



Artesunate Injection IP

Artecan-120

120 mg Injection

Artesunate Injection (Artesunate 120 mg / Vial)

COMPOSITIONS:

Each vial contains 120mg of Artesunate Injection IP

Each ampoule contains 2 ml of 5% Sodium bicarbonate solution.

1 ampoule of 10 ml ampoule of Sodium Chloride Injection IP 0.9% w/v

ACTIONS:

- Artesunate is an antimalarial agents. Artesunate is a derivative of Artemisinin found in *Artemisia annua*, a herb that has traditionally been used in China for the treatment of malaria.
- Artesunate is an alternative therapy for the treatment of severe malaria, including cerebral malaria.

INDICATIONS:

Artesunate has an antimalarial activity. It can quickly and reliably control the acute attack and cerebral type of malaria. It is effective against *plasmodium falciparum* and *plasmodium vivax*, especially against malaria parasites resistant to Chloroquine.

ADMINISTRATION:

- As prescribed by physicians.
- It can be given by deep IM injection or slow IV injection:
 - + Deep IM injection: Dissolve 120mg of Artesunate with 2ml of 5% sodium bicarbonate solution for injection before use.
 - + Slow IV injection: Dissolve 120mg of Artesunate with 2ml of 5% sodium bicarbonate solution for injection and add 10ml of 0.9% sodium chloride solution for injection before use to make 1ml contains 10mg of Artesunate. Injection must follow immediately soon after dissolution, if the solution appears cloudy or sediment occurs, it should be rejected.

DOSAGE:

The usual injection dosage for each time:

- Adult and Children > 7 years: 1.2mg/kg.

The following doses are given at hour 4 and 24 after the first injection in day 1. Give 1 dose daily for the consecutive days. A course of therapy is recommended for 5 consecutive days.

CONTRA – INDICATIONS:

No contra-indications has been so far reported.

SIDE-EFFECTS:

The most frequent side-effect are mild, transient such as: side-effects on gastrointestinal system (nausea, abdominal pain, diarrhea), headache, dizziness.

PRECAUTIONS:

Use with caution in pregnant women in the first trimester.

INTERACTIONS:

Due to its structure containing a sesquiterpene lactone that bears a peroxide grouping which releases free oxygen, avoid using Artesunate concomitantly with antioxidants such as Vitamin E, Vitamin C, Glutathione, because those will decrease the antimalarial action of the drug.

SHELF-LIFE:

24 months from the manufacturing date.

Never use after the expiry date clearly indicated on the outer packaging.

STORAGE: Store in a cool, dry place. Protected from direct sunlight. Do not freeze.

Improper storage may deteriorate the product.

PRESENTATION:

Each box contains:

- 1 vial of Artesunate Injection IP 120 mg
- 1 ampoule of 2ml of 5% Sodium bicarbonate Injection IP.
- 1 ampoule of 10 ml ampoule of Sodium Chloride Injection IP 0.9% w/v

Manufactured by:



**AMERICAN
REMEDIES**[®]
Care & Cure with Research

American Remedies Healthcare Pvt. Ltd.
(An ISO 9001-2015 Certified Company)

ICO: American Remedies LLC, DE, USA



Gala No. 105 to 111, 1st Flr., Bldg. No. E-15/A,
Harihar Complex, Bhiwandi, Mumbai - 421302

At: Vill. Surajpur Nahar Road Paonta Sahib,
Distt. Sirmour, Himachal Pradesh-173001

E-mail: info@americanremedies.in

© Copyright

Scan for
Batch
Certificate

