

For the use of a Registered Medical Practitioner OR A Hospital OR A Laboratory.


Citicoline Sodium Injection IP

AMERICAN Citistroke 4 Injection

Composition :
Each ml contains :
Citicoline Sodium IP
eq. to Citicoline 250 mg
Water for Injections IP q.s.

For I.M./I.V. Use Only

Description:
Citicoline is a complex organic molecule that functions as an intermediate in the biosynthesis of cell membrane phospholipids. Citicoline is also known as CDP-choline and cytidinediphosphate choline (cytidine 5'-diphosphocholine). CDP-choline belongs to the group of biomolecules in living systems known as "nucleotides" that play important roles in cellular metabolism. CDP-choline is composed of ribose, pyrophosphate, cytosine (a nitrogenous base), and choline. Exogenous citicoline research in animal experiments and human clinical trials provides evidence of its cholinergic and neuroprotective actions. As a dietary supplement, citicoline appears useful for improving both the structural integrity and functionality of the neuronal membrane that may assist in membrane repair.



Therapeutic Indications
For the treatment of disturbances of consciousness as resulting from head injuries, brain operation and acute stage of cerebral infraction in adult only.
Posology and Method of Administration
Disturbance of Consciousness resulting from head injury or brain operation
Usually for adults a dose of 500-1000 mg of Citistroke-4 Injection is administered once or twice a day, by intravenous drip, infusion, intravenous injection or intramuscular injection. The dose may be adjusted according to the patient's age and condition.
Disturbance of Consciousness in the Acute Stage of Cerebral Infarction
Usually, a dose of 1,000 mg of citicoline is administered once a day, by intravenous injection, for 2 consecutive weeks.
The dosage may be adjusted based on the seriousness of the disease. It can be administered intramuscularly, intravenously (3 to 5 minute) injection and in intravenous drop perfusion (dripping speed 40-60 drops/minute). Citicoline is compatible with all intravenous isotonic solutions. It can also be mixed with hypertonic glucose serum.
Contraindications
It is contraindicated in patients with a known hypersensitivity to citicoline or any other component of the formulation. Citicoline must not be administered to patients with hypertonic of the parasympathetic.
Special Warnings and Precautions for Use
Cholines are generally regarded as safe and appear to be well-tolerated. High intake of cholines may cause low blood pressure, steatorrhea (undigested fat in stool), nausea, vomiting, salivation, diarrhoea, constipation, anorexia, dizziness (vertigo), sweating, insomnia and headache. Cholines can possibly trigger existing epilepsy. In administering citicoline injection intramuscularly, caution should be exercised so as not to affect the tissues, nerves, etc. Intramuscular injection should be given only when indispensable and should be restricted to the minimum to be required. In particular, repeated injection at the same site should be avoided. Care should be exercised to avoid injection at sites along the course of the nerves. In case of intense pain or backflow of blood upon insertion of the injection needle, the needle should be withdrawn immediately and injected at a different site. In intravenous administration, inject as slowly as possible. Since shock may occur, a close observation should be maintained. If any such abnormalities such as drop in blood pressure, distressed feeling of the chest or dyspnoea are observed, citicoline injection should be discontinued and appropriate measures taken.
Drug Interactions
Citicoline may enhance the effects of levodopa. The exact mechanism is unknown, but animal models suggest that citicoline may increase dopamine levels in the brain and/or improve dopaminergic cell survival. In patients with Parkinson's disease, a few studies have demonstrated levodopa-saving effects, whereby the addition of citicoline (500 to 1200 mg/day) allowed for lower dosages of levodopa to be used with stable or improved therapeutic efficacy and reduced side effects in some patients. However, data are limited.

Coadministration with meclofenoxate
Citicoline must not be administered in conjunction with medication containing meclofenoxate (also known as clophenoxate).
Use in Special Population
Patients with Renal Impairment
Adjustment of the dose is recommended in elderly patients with compromised renal function. For long-term treatment in the elderly, regular evaluation of the creatinine clearance is required to allow dosage adaptation if needed. The daily dose must be individualized according to renal function.
Patients with Hepatic Impairment
There are no data concerning the effects of liver insufficiency on the safety profile and pharmacokinetics of citicoline.
Pregnant Women
There are no adequate and well controlled studies of citicoline during pregnancy and lactation. Citicoline should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus.
Lactating Women
There are no adequate and well controlled studies of citicoline during pregnancy and lactation. Caution should be exercised during breastfeeding because it is not known whether citicoline is excreted in human breast milk.
Paediatric Patients
The experience in children is limited; therefore, it may only be administered when the expected therapeutically benefit is higher than any possible risk.
Geriatric Patients
No dosage adjustment is required in this patient population and the usually recommended adult dose can be administered. However, adjustment of the dose is recommended in elderly patients with compromised renal function.
Effects on Ability to Drive and Use Machines
Patient are advised not to operate heavy machinery or automobiles until the full effect of citicoline is known.
Undesirable Effects
The commonly observed adverse effects (0.1–5%) with intravenous use of citicoline were rash, insomnia, occurrence or intensification of numbness or paralyzed extremities (when used in patients with post-apoplectic hemiplegia), nausea, abnormal laboratory values for function of the liver, and feeling of warmth. The other adverse reactions (<0.1%) were excitation, convulsions, anorexia, transient diplopia, transient blood pressure changes, malaise, shock, distressed feeling of the chest, and dyspnea. In a short-term, placebo-controlled, crossover study, 12 healthy adults took citicoline at daily doses of 600 mg and 1,000 mg or placebo for consecutive 5-day periods. Transient headaches occurred in 4 subjects on the 600 mg dose, 5 on the 1,000 mg

dose, and 1 on placebo. No changes or abnormalities were observed in hematology, clinical biochemistry or neurological tests. A large drug surveillance study analyzed the results of citicoline treatment in 2,817 patients aged 60 to 80 years, suffering from senility and cerebral vascular insufficiency. A total of 151 incidents of side effects were recorded, representing 5% of the patient sample. The most common adverse effects were transient in nature and included stomach pain and diarrhoea in 102 cases. Vascular symptoms of hypotension, tachycardia or bradycardia occurred in 16 cases.
Citicoline is generally well tolerated. Few adverse effects that are reported with oral citicoline include gastrointestinal disturbances, dizziness and fatigue, rash, insomnia, nausea, dizziness, convulsions, anorexia, shock, transient blood pressure changes.

Overdose
Citicoline exhibits very low toxicity profile in humans. In a short term, placebo controlled, crossover study, 12 healthy adults took citicoline at daily doses of 600 and 1000 mg or placebo consecutive 5-days periods. Transient headaches occurred in 4 subjects on 600 mg dose, 5 on the 1000 mg dose and 1 in placebo. No changes or abnormalities were observed.
Pharmacological Properties
Mechanism of Action
Citicoline acts by various mechanisms as cerebral activator listed below:
Phospholipid Precursor
Evidence of citicoline's role as a phosphatidylcholine precursor has been found in animal studies. The brain uses choline preferentially for acetylcholine synthesis, which can limit the amount of choline available for phosphatidylcholine production. When the demand for acetylcholine increases or choline stores in the brain are low, phospholipids in the neuronal membrane can be catabolized to supply the needed choline. Exogenous citicoline thus helps preserve the structural and functional integrity of the neuronal membrane. In an *in vitro* study, citicoline at high concentrations stimulated brain acetylcholinesterase (AChE) along with Na⁺/K⁺-ATPase. The postulated mechanism involves bioconversion of citicoline to phosphatidylcholine.
Neuronal Membrane Repair
Citicoline has been investigated as a therapy for stroke patients. Three mechanisms are postulated: (1) repair of neuronal membranes via increased synthesis of phosphatidylcholine; (2) repair of damaged cholinergic neurons via potentiation of acetylcholine production; and (3) reduction of free fatty acid buildup at the site of stroke-induced nerve damage. In addition to phosphatidylcholine, citicoline serves as an intermediate in the synthesis of sphingomyelin, another neuronal membrane phospholipid component. Citicoline has shown the potential to restore post-ischemic sphingomyelin levels.
Citicoline also restores levels of cardiolipin, a phospholipid component of the inner mitochondrial membrane. The mechanism for this is unknown, but data suggest citicoline inhibits enzymatic hydrolysis of cardiolipin by phospholipase. In an animal study, citicoline decreased the formation of hydroxyl radicals following ischemia and perfusion, again suggesting citicoline acts to decrease phospholipase stimulation.

Effect on beta-Amyloid
Evidence has surfaced that citicoline counteracts the deposition of beta-amyloid, a neurotoxic protein believed to play a central role in the pathophysiology of Alzheimer's disease (AD). The characteristic lesion in AD is the formation of plaques and neurofibrillary tangles in the hippocampus. The degree of cognitive dysfunction and neurodegeneration in AD is proportional to the buildup of beta-amyloid. Citicoline counteracted neuronal degeneration in the rat hippocampus induced by intra hippocampal injection of beta-amyloid protein. The number of apoptotic cells was also reduced. Memory retention as measured by a passive-avoidance learning task improved in the rats.
Effect on Neurotransmitters
Evidence of citicoline's ability to enhance norepinephrine release in humans was found in a study showing citicoline raised urinary levels of 3-methoxy- 4-hydroxyphenylglycol (MHPG), a norepinephrine metabolite. Citicoline increased brain levels of neurotransmitters in rats at a dose of 100 mg/kg, administered daily for seven days. Norepinephrine increased in the cerebral cortex and hypothalamus; dopamine increased in the corpus striatum, and serotonin increased in the cerebral cortex, striatum, and hypothalamus. Rat studies have found evidence that citicoline potentiates dopamine release in the brain, presumably by stimulating release of acetylcholine.

Pharmacodynamic Properties
When administered orally, it is absorbed almost completely, and its bioavailability is approximately the same when administered intravenously. Once absorbed, the cytidine and choline disperse widely throughout the body, cross the blood-brain barrier, and reach the central nervous system (CNS), where they are incorporated into the phospholipids fraction of the cellular membrane and microsomes. The concept that the administration of exogenous Citicoline can augment the synthesis of neural membrane phospholipids is attractive, because accelerated replacement or repair plays a critical role in maintaining the healthy function of numerous physiological processes. It has shown the therapeutic efficacy in a variety of diseases in which membrane disorder, dysfunction, or degeneration result in cellular and tissue ischaemia and necrosis.

Pharmacokinetic Properties
Citicoline is well absorbed following intramuscular administration. After intramuscular doses of citicoline 1,000 mg, peak increases in plasma choline levels were seen in 0.4 hours, with levels increasing from 11 micromol/L (baseline) to 25 micromol/L.
Choline derived from citicoline crosses the blood-brain barrier, presumably serving as a source of acetylcholine and phosphatidylcholine (lecithin) synthesis. The major portion of a dose of citicoline appears to be incorporated into tissues and/or used in biosynthetic/ biodegradation pathways, including lecithin/lipid membrane synthesis.
Citicoline is metabolized in the liver to free choline. The liver is capable of synthesizing lecithin from choline. The half-life of free choline is of 2 hours after intramuscular administration.
Only small amounts of dose are recovered in the urine and faeces (less than 3% each). Approximately 12% of a dose is eliminated through the lungs as carbon dioxide.

Nonclinical Properties
Animal Toxicology or Pharmacology
Acute toxicity from single citicoline administration has been studied in various animal species and using different administration routes. The intravenous LD50 in mice, rats, and rabbits is 4.6, 4.15, and 1.95 g/kg, respectively. Oral LD50 is 27.14 g/kg in mice and 18.5 g/kg in rats. The intravenous LD50 of citicoline is approximately 44 times higher than the LD50 of choline hydrochloride at equivalent doses, and it has been shown that choline doses inducing cholinergic crises do not cause any toxicity sign when equivalent doses of citicoline are administered. This suggests that administration of choline has metabolic implications clearly different from those of exogenous choline administration. The administration of 2000 mg/kg of citicoline during 14 days was well tolerated. Intraperitoneal administration to rats of doses up to 2 g/kg/d of citicoline for 4.5 weeks did not result in clinical toxicity signs or significant changes in the haematological, biochemical, or histological parameters analysed. A slight decrease in intake and weight gain was only seen from 2 weeks of the study. Similar results were seen following subcutaneous administration to male rats of up to 1 g/kg for 4 weeks. Oral administration of 1.5 g/kg/d to rats for 30 days did not cause weight, haematological, biochemical, or histological changes.
Chronic oral (1.5 g/kg/d for 6 months in dogs) and intraperitoneal (1 g/kg/d for 12 weeks in rats) toxicity studies did not reveal either significant abnormalities related to drug administration. Intravenous administration of citicoline 300-500mg/kg/d for 3 months in dogs only caused toxic signs immediately after injection, including vomiting and occasional diarrhoea and sialorrhoea. In a 90-day study in rats, 100, 350, and 1000 mg/kg/day oral doses resulted in no mortality. In males, slight significant increases in serum creatinine (350 and 1000 mg/kg/day) and decreases in urine volume (all treated groups) were observed. In females, slight significant increases in total white blood cell and absolute lymphocyte counts (1000 mg/kg/day), and blood urea nitrogen (BUN) (100 and 350, but not 1000 mg/kg/day) were noted. A dose-related increase in renal tubular mineralization, without degenerative or inflammatory reaction, was found in females (all treated groups) and two males (1000 mg/kg/day). Renal mineralization in rats (especially females) is influenced by calcium: phosphorus ratios in the diet. A high level of citicoline consumption resulted in increased phosphorus intake in the rats, and likely explains this result.

Storage: Store protected from light, at a temperature not exceeding 25°C.

Marketed by:



Gala No. 105 to 101, 1st Flr., Bldg. No. E-15/A,
Haitari Complex, Bhiwandi, Mumbai - 421302
E-mail: info@americanremedies.in
ICO: American Remedies LLC, DE, USA
© Copyright