

Holybion Injection

Vitamin B Complex Injection

Composition:

Each ml contains	
Thiamine HCl	I.P. 5mg
Riboflavin sodium phosphate	I.P. 2mg
Cyanocobalamin	I.P. 4mg
Niacinamide	I.P. 20mg
Pyridoxine HCl	I.P. 2mg
D-Panthenol	I.P. 3mcg
Benzyl Alcohol	I.P. 0.9%V/V
(As preservative)	
Water for injection	I.P. q.s

Indication:

Vitamin B complex is indicated in following diseases: vitamin B deficiency and (neuritis); alcohol, toxic, and post infections polyneuritis; diabetic polineuropathy; neuralgia; sciatica; central spastic conditions; myasthenia; paresthesia; atheroscleroses; Wernicke's encephalopathy; vegetative neuroses; dermatitis; neurodermatitis; psoriasis; lupus erythematoses; furunculosis; stomatitis; cheilitis; glossitis; colitis; hepatitis; chronic alcohol

Dosage and Administration:

It is usually given 0.25 to 2ml by intramuscular or slow intravenous injection. High concentrations given intravenously may be diluted using parenteral infusion solutions.

Contraindications:

Hypersensitivity to the active substances or to any of the excipients

Warnings and precautions:

Anaphylactogenesis may occur with parenteral thiamine. Use with caution. An intradermal test dose is recommended prior to administration in patients suspected of being sensitive to the drug.

Pregnancy and Lactation:

Vitamins from B group are generally recommended during pregnancy and breastfeeding as per individual requirement. These vitamins pass through breast milk to the infant. No specific safety data could be identified. Physician's advice before use is recommended.

Drug Interactions:

Some medication may interact with vitamin B complex. Before using this product, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: altretamine, cisplatin certain antibiotics (e.g., chloramphenicol), certain anti-seizure drugs (e.g., phenytoin), levodopa, other vitamin/nutritional supplements. Consult with your doctor about any medications you are taking, before your treatment with vitamin B complex injection.

Adverse Effects:

Mild transient diarrhea, polycythemia vera, peripheral vascular thrombosis, itching transitory exanthema, feeling of swelling of entire body; anaphylactic shock and death. Use should be discontinued upon observance of any untoward reaction. Pain upon intramuscular injection may be noted.

Overdose:

Symptoms of overdose include thirst, flushing of the skin, and abdominal pain and dizziness, redness of the skin, excessive urination and diarrhea. These symptoms have to be reported to the medical health professional. Seek medical help urgently.

Storage : Store in a cool, dry and dark place. Improper storage may deteriorate the product.

Packing : 10x5x2ml

Marketed by:



Care & Cure with Research
American Remedies Healthcare Pvt. Ltd.

(An ISO 9001-2015 Certified Company)

ICO: American Remedies LLC, DE, USA

Gala No. 105 to 111, 1st Flr., Bldg. No. E-15/A,
Harihar Complex, Bhiwandi, Mumbai - 421302

E-mail: info@americanremedies.in

© Copyright