

For the use of registered medical practitioner or a Pharmacist or a Laboratory

Metoclopramide Hydrochloride Injection, I.P.

Composition :
Each 2ml contains :
Metoclopramide Hydrochloride IP
Equivalent to anhydrous
Metoclopramide Hydrochloride 10mg
Benzyl Alcohol IP 2% w/v
(As preservative)
Water for injection IP q.s.

1. Name of the medicinal Product
Metoclopramide Injection IP 10 mg/2 ml

2. Qualitative and Quantitative Composition
Each 2 ml contains metoclopramide hydrochloride IP equivalent to 10 mg of anhydrous metoclopramide hydrochloride

3. Pharmaceutical Form
Solution for Injection
A clear colorless solution.

4. Clinical Particulars

4.1 Therapeutic Indications Adults (20 years and over):
As an adjunct to X-ray examination of the stomach and duodenum. To assist in intestinal intubation and control nausea and vomiting associated with the following conditions: intolerance to essential drugs possessing emetic properties, uraemia, radiation sickness, malignant disease, postoperative vomiting, and lab. or infectious diseases. There is no clear benefit in motion sickness or other labyrinth disturbances. I.M administration of **METOCAN** injection facilitates the absorption of a range of drugs including the absorption of aspirin in people with migraine. **METOCAN** injection may found useful in the management of gastric retention after gastric surgery. **METOCAN** injection may be useful in the treatment of diabetic gastroparesis of mild to moderate severity. Once control of diabetes has been established by diet and/or insulin, **METOCAN** injection should be avoided.

Young adults and children over 1 year of age: The use of **METOCAN** injection in patients under 20 years should be restricted to the following situations and only used as second line therapy. Severe intractable vomiting of known cause. Vomiting associated with radiotherapy and intolerance to cytotoxic drugs. As an aid to gastrointestinal intubation.

4.2 Posology and Method of Administration
Route of administration: IM or by slow IV injection (1 to 2 minutes)
Patients with normal renal and hepatic function: The dosage recommendations given below should be strictly adhered to if side effects of the dystonic type are to be avoided. Total daily dosage of **METOCAN** injection, especially for children and young adults should not normally exceed 0.5 mg/kg bodyweight with a maximum of 30 mg daily. **METOCAN** injection should only be used after careful examination to avoid masking an underlying disorder, e.g., cerebral irritation. Maximum recommended treatment duration is 5 days in all age groups. Adults 20 years and over: Maximum of 20 mg three times a day, 1 to 3 times daily, IV or I.M. depending on the severity of the condition. Elderly patients: To avoid adverse reactions strict adherence to dosage recommendations is advised and, where prolonged therapy is considered necessary, patients should be regularly reviewed.

Young adults and children over 1 year of age: Treatment of children should commence at the lower dosage, where advised, and used as second line therapy only.
Young adults: 15-19 years: Treatment of young adults should commence at the lower dosage, and used as second line therapy only 5 mg to 10 mg three times a day. Children 5 - 14 years: 2.5 mg to 5 mg (0.5 mL to 1 mL) two or three times daily 3-5 years: 2 mg two or three times daily. Children 1 - 3 years: 1.0 mg two or three times daily.

Diagnostic indications: A single dose of may be given 5 to 10 minutes before the examination. Subject to body weight considerations, the following dosages are recommended:
Adults: 20 years and over: 10 to 20 mg
Young adults: 15-19 years: 10 mg
Children: 5-14 years: 5 mg, 5-9 years: 2.5 mg, 3-5 years: 2 mg, 1-3 years: 1 mg.
Patients with impaired renal and hepatic function: In patients with clinically significant degrees of renal or hepatic impairment, clearance of **METOCAN** injection is likely to be reduced. It is suggested that therapy be initiated at half the recommended dose. Subsequent dosage will depend on individual clinical response.

4.3 Contraindications
It should not be used whenever stimulation of gastrointestinal motility might be dangerous, e.g., in the presence of gastrointestinal haemorrhage, mechanical obstruction, or perforation.
4.4 Special Warnings and Special Precautions for Use
Intravenous administration of **METOCAN** injection should be given slowly over a 1 to 2 minute period, since a transient but intense feeling of anxiety and restlessness, followed by drowsiness, may occur with rapid administration. Persistent tardive dyskinesia: Treatment with **METOCAN** injection can cause tardive dyskinesia (TD), a potentially irreversible and disabling disorder characterized by involuntary movements of the face, tongue, or extremities. Treatment with **METOCAN** injection for longer than the recommended 12 weeks should be avoided in all but rare cases where therapeutic benefit is thought to outweigh the risk of developing TD.

METOCAN injection should be discontinued in patients who develop signs or symptoms of TD. There is no known effective treatment for established cases of TD. Therefore, **METOCAN** injection should not be used for the symptomatic control of TD. Extrapyramidal Symptoms (EPS) may occur with both **METOCAN** injection and neuroleptics such as phenothiazines; care should be taken in the event of both drugs being prescribed concurrently. **METOCAN** injection given to epileptic patients the frequency and severity of seizures or extrapyramidal reactions may be increased. Acute dystonic reactions occur in approximately 1% of patients given **METOCAN** injection. These occur more frequently in children and young adults and may occur after a single dose. **METOCAN** injection monotherapy may report neuroleptic malignant syndrome, combination with neuroleptics. Symptoms: hyperpyrexia, altered consciousness, muscle rigidity, autonomic instability and elevated levels of CPK and must be treated urgently (measured treatments include dantrolene and bromocriptine). The treatment should be avoid immediately if this syndrome occurs.

Protein levels: **METOCAN** injection elevates prolactin levels and the elevation persists during chronic administration. A factor of potential importance if the prescription of metoclopramide is contemplated in a patient with previously detected breast cancer. Although disturbances such as galactorrhoea, amenorrhoea, gynaecomastia, and impotence may be reported with prolactin elevating drugs.
Other: **METOCAN** injection used (in such operation, pyloroplasty or gut anastomosis) should be withheld for three or four days as vigorous muscular contractions may not help healing. The symptomatic relief provided by **METOCAN** may delay recognition of serious disease. It should not be prescribed until diagnosis has been established and should not be substituted for appropriate investigation of the patient's symptoms.
If vomiting persists in a patient receiving **METOCAN** injection, the patient should be reassessed to exclude the possibility of an underlying disorder (e.g. cerebral irritation). **METOCAN** injection may induced depression in patients without a prior history of depression. It should be given to patients with a prior history of depression only if the expected benefits outweigh the potential risks.
METOCAN injection should be used with caution in patients with hypertension as on (I.V) administration may show to release catecholamines.
Elderly: To avoid adverse reactions adhere strictly to dosage recommendations and where prolonged therapy is considered necessary, patients should be regularly monitored.
Pediatric use: It is contraindicated in children less than 1 year of age thus it should not be used in this age group. Unless a clearly indication has been established for its use, because of the to avoid higher incidence of adverse reactions.
Pregnancy: No adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if clearly needed.
Lactation: Metoclopramide is excreted in human milk. Caution should be exercised when metoclopramide is administered to a nursing mother.

4.5 Interaction with other medicinal products and other forms of interaction
The effects of metoclopramide on gastrointestinal motility are antagonized by anticholinergic drugs and narcotic analgesics. Additive sedative effects can occur when metoclopramide is given with alcohol, sedatives, hypnotics, narcotics, or tranquilizers. Since metoclopramide accelerates abnormally slow gastric and small bowel peristaltic activity. The absorption of drugs from the small bowel may be accelerated (e.g. paracetamol, tetracycline, L-dopa), whereas absorption of drugs from the stomach may be diminished (e.g. digoxin). Metoclopramide may cause extrapyramidal symptoms in some patients. Therefore, when metoclopramide is used concomitantly with other drugs that are likely to cause extrapyramidal reactions, (e.g., neuroleptics such as phenothiazines), caution should be taken. The decrease in gastric emptying time caused by metoclopramide may increase the bioavailability of cyclosporin. Monitoring of cyclosporin concentrations may be necessary. When metoclopramide is given concurrently with sucralfate, the recovery time is prolonged. Metoclopramide influences the delivery of food to the intestine and thus, the rate of its absorption, the administration of metoclopramide may result in poor diabetic control in some patients. Therefore, adjustment in, or timing of, insulin dosage may be necessary in insulin controlled diabetics.

The finding that metoclopramide releases catecholamines in patients with essential hypertension suggests that it should be used cautiously, if at all, in patients receiving monoamine oxidase inhibitors.
4.6 Fertility, Pregnancy and Lactation
Pregnancy: No adequate and well-controlled studies in pregnant women. This drug should be used during pregnancy only if clearly needed.
Lactation: Metoclopramide is excreted in human milk. Caution should be exercised when metoclopramide is administered to a nursing mother.

4.7 Effects on ability To Drive and use Machines.
Patients should be cautioned about engaging in activities requiring mental alertness for a few hours after the drug has been administered.

4.8 Undesirable Effects
Less frequently, insomnia, headache, dizziness, nausea, or bowel disturbances may occur. Rarely may occur acute depression.
Anxiety or agitation also may occur, especially after rapid injection.
A single instance of supraventricular tachycardia following (IM) administration may be report.

There may very rare of abnormalities of cardiac conduction (such as bradycardia and heart block) in association with intravenous metoclopramide. Atrial fibrillation, oedema, tachycardia and palpitations may be associated with the use of metoclopramide. Extrapyramidal Reactions, Symptoms include: spasm of the facial muscles, trismus, rhythmic protrusion of the tongue, a bulbar type of speech, spasm of the extra ocular muscles including oculogyric crises, unusual positioning of the head and shoulders and opisthotonus. There may be a generalized increase in muscle tone. The majority of reactions occur within 36 hours of starting treatment and the effects usually disappear within 24 hours of withdrawal of the drug, an antiparkinson drug such as benzhexol or an anticholinergic antihistamine such as diphenhydramine should be given. A fatal dystonic reaction may report in a patient who received hexamethylmelamine, cisplatin and high dose metoclopramide. A fatal cardiorespiratory arrest may occur in at least one patient with an acute dystonic reaction.

Tardive dyskinesia, which may be persistent, may report particularly in elderly patients undergoing long-term therapy with **METOCAN** injection.
Neuroleptic Malignant Syndrome: Very rare cases occurrences of this syndrome is potentially fatal and comprises hyperpyrexia, altered consciousness, muscle rigidity, autonomic instability and elevated levels of CPK and must be treated urgently (measured treatments include dantrolene and bromocriptine). The treatment should be avoid immediately if this syndrome occurs.
Parkinsonian symptoms, including tremor, rigidity, bradykinesia and akinesia, occur rarely in patients Treatment with metoclopramide but may be associated with usual or excessive doses or with decreased renal function.

There may causes of hypersensitivity reactions such as urticaria, maculopapular rash, a limited case of neutropenia, leucopenia, Sulphamoglobinaemia (in adults), methaemoglobinemia may also report and agranulocytosis generally without perfect association with metoclopramide.
Galactorrhoea and breast enlargement, raised serum prolactin levels.
Respiratory failure, secondary to dystonic reaction, acute asthmatic symptoms of wheezing and dyspnoea may occur. Urinary incontinence and frequency, sexual dysfunction, priapism and muscle spasm may also occur. Rarely, cases of hepatotoxicity characterized by such as jaundice and altered liver function tests, when metoclopramide taken with other drugs with known hepatotoxic potential.

4.9 Overdose
Symptoms: The overdose of metoclopramide can rise, Drowsiness, disorientation and extrapyramidal, precise hardy AV block, feelings of anxiety or restlessness, headache, vertigo, nausea, vomiting, constipation, weakness, hypotension and vasodilation.
Treatment: The overdose of metoclopramide symptomatic and supportive therapy may be followed. Antiparkinson and anticholinergic drugs such as diphenhydramine hydrochloride have effectively controlled extrapyramidal reactions. Haemodialysis appears ineffective in removing metoclopramide. Similarly, continuous ambulatory peritoneal dialysis does not remove significant amounts of the drug.

5. Pharmacological Properties
5.1 Pharmacodynamics Properties
Metoclopramide has antiemetic, antinauseant, and gastrokinetic activity. It stimulates motility of the upper gastrointestinal tract without stimulating gastric, biliary or pancreatic secretions. The rate of gastric emptying is increased due to increased peristalsis of the jejunum and duodenum.
The tone and amplitude of gastric contractions are increased, with relaxation of the pyloric sphincter and duodenal bulb. These effects combine to result in decreased intestinal transit time. The effect of metoclopramide on motility is not dependent on intact vagal innervation, but it can be abolished by anticholinergic drugs. Metoclopramide has little, if any, effect on the motility of the colon or bladder. Metoclopramide also exhibits dopamine antagonist activity and consequently produces sedation and, rarely, other extra pyramidal reactions. It may have serotonin receptor (5HT₂) antagonist properties. Metoclopramide inhibits the central and peripheral effects of apomorphine, induces release of prolactin and produces a transient increase in circulating aldosterone levels.

5.2 Pharmacokinetic Properties
Absorption: Following intravenous administration, the onset of action is within 1 to 3 minutes and after intramuscular administration, this interval is extended to 10 to 15 minutes. The effect usually lasts from 1 to 2 hours.
Protein Binding: Plasma protein binding is 13% to 22%.
Metabolism: About 80% of the drug is excreted in the urine in the first 24 hours after administration. Approximately half is unchanged metoclopramide and half is the glucuronide and sulphate conjugate.
Elimination: Metabolism mainly occurs in the liver and elimination half-life may vary from 2.5 to 6 hours. Impaired renal function results in a reduced clearance and an increased half-life, up to 15 hours. The drug is not extensively bound to plasma proteins. The whole body volume of distribution is high which suggests extensive distribution of drug to the tissues.

5.3 Preclinical Safety Data
No data available.
5.4 Pharmaceutical Particulars

5.1 Shelf Life
24 months
5.2 Special Precautions for Storage
Storage - Store protected from light.
5.3 Special precautions for disposal and other handling
Any unused product or waste material should be disposed of in accordance with local requirements.

5.4 Shelf Life
24 months
5.2 Special Precautions for Storage
Storage - Store protected from light.
5.3 Special precautions for disposal and other handling
Any unused product or waste material should be disposed of in accordance with local requirements.

5.4 Shelf Life
24 months
5.2 Special Precautions for Storage
Storage - Store protected from light.
5.3 Special precautions for disposal and other handling
Any unused product or waste material should be disposed of in accordance with local requirements.

Marketed by:
AMERICAN REMEDIES
Care & Cure with Research
American Remedies Healthcare Pvt. Ltd.
(An ISO 9001:2015 Certified Company)
Gala No. 105 to 111, 1st Flr., Bldg. No. E-15/A, Harihar Complex, Bhiwandi, Mumbai - 421302
E-mail: info@americanremedies.in
ICO: American Remedies LLC, DE, USA
© Copyright

