

For The Medical Practitioner Or A Laboratory Use Only

Lignocaine Hydrochloride Gel IP

NumbOn 2% Gel

with applicator

Composition

Lignocaine Hydrochloride IP equivalent to	
Lignocaine Hydrochloride Anhydrous	2.000% w/v
Methylparaben IP	0.061% w/v
Propylparaben IP	0.027% w/v
(as preservatives)	
Water soluble gel base	q.s.

PHARMACEUTICAL FORM:

Gel Form

INDICATIONS:

This medication is used to prevent and control pain during certain medical procedure such as inserting a tube into the urinary track. It is also used to numb the lining of the mouth, nose, throat, or urinary tract (e.g., endotracheal intubation, urinary catheterization). Lignocaine gel is also used to numb and treat inflammation of the urinary tract (urethritis).

CONTRA-INDICATION:

Traumatized mucosa, secondary bacterial infection of the area of proposed application and known hypersensitivity to any of the components. Lignocaine Hydrochloride Gel IP is contraindicated in patients with a known history of hypersensitivity to local anesthetics of the amide type.

DOSAGE AND ROUTE OF ADMINISTRATION:

Cleanse the wound and blow it dry. Apply a thin layer of Lignocaine Hydrochloride Gel IP 2%w/v to the wound surface and the skin immediately surrounding the wound 3-4 times daily.

Route of Administration: through Topical administration only. (For external use only.)

OVER DOSE:

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, consult with physician immediately.

ADVERSE EFFECTS:

Adverse experiences following the administration of Lignocaine Hydrochloride Gel IP are similar in nature to those observed with other amide local anesthetic agents. These adverse experiences are, in general, dose-related and may result from high plasma levels caused by excessive dosage or rapid absorption, or may result from a hypersensitivity, idiosyncrasy or diminished tolerance on the part of the patient.

Serious adverse experiences are generally systemic in nature. The following types are those most commonly reported.

SPECIAL PRECAUTION:

Do not use this product if you are allergic to any ingredients. If condition worsens or does not improve within 7 days, consult a physician. Do not use on children under 2 years of age without consulting a physician.

Avoid contact with eyes. Do not use in large quantities.

DRUG INTERACTION:

Consult your physician if any see the reaction on the skin after using this medicine.

PREGNANCY AND LACTATION:

Pregnancy:

Consult your physician before administering this medicine during pregnancy.

Lactation:

Consult your physician before administering this medicine during lactation.

WARNING: Keep out of reach children.

STORAGE: Store at a temperature not exceeding 30°C. The Gel should not be frozen.

PRESENTATION: 1 x 30 g Aluminum Tube + Applicator In Blister Pack.

Marketed by:

 **AMERICAN
REMEDIES**
Care & Cure with Research
American Remedies Healthcare Pvt. Ltd.
(An ISO 9001-2015 Certified Company)

Gala No. 105 to 111, 1st Flr., Bldg. No. E-15/A,
Harihar Complex, Bhiwandi, Mumbai - 421302
E-mail: info@americanremedies.in
ICO: American Remedies LLC, DE, USA 
© Copyright

Scan for
Batch Certificate

