

160mm

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only.

Acetaminophen Suppositories USP 170 mg (Paracetamol Suppositories)

AMERICAN
PARACAN 170
Suppositories

COMPOSITION:
Each Suppository contains:
Paracetamol IP170 mg.
Suppository Baseq.s.
Colour: Titanium Dioxide.

INDICATIONS:
Symptomatic relief of pain and fever. For children below 6 years.

CONTRA-INDICATIONS:
This medicine must not be used in patients with: known hypersensitivity towards the active ingredient Paracetamol Infants from birth up to the end of the second month. Special care should be taken when using this medicine (That is, with a prolonged interval between doses or at a diminished dose) and under medical supervision inpatients with: Disorders of liver function (e.g. chronic alcohol misuse, inflammatory conditions of the liver), Disorders of renal function.

ADVERSE REACTIONS:
Reddening of the skin may occur, through rarely; allergic reactions with cutaneous eruptions are very rare.

DRUG INTERACTION:
The medicine contains Paracetamol. It should not be continued with other Paracetamol preparations so as not to exceed the recommended daily dose. Drugs which induce hepatic microsomal enzymes such as Alcohol, Rifampicin, Barbiturates and other anti-convulsions may increase the hepatotoxicity of Paracetamol particularly after overdosage. Therapeutic effects of Paracetamol may also be decreased.

DOSAGE AND ADMINISTRATION:
The dosage of Paracetamol is governed by age and body weight, and is usually 10-15 mg Paracetamol per kg body weight as single dose and upto 50 mg/ kg body as a maximum dose per day (24 hours). The recommended usual daily dose depends on the weight of the infant.

Dosage Instruction
For information only

Baby (till 6 months) up to 7 kg: 80 mg suppository, 4times daily
Infants (1 to 3 years)/from 10 to 12 kg: 125 mg Suppository 4 times daily
Children (2 to 4 years)/from 12 to 16 kg: 170 mg Suppository 4 times daily
Children (4 to 7 years)/from 17 to 22 kg: 250 mg Suppository 4 times daily

The approximate age as a function of body weight are given for example only. In case of any doubt, the doctor should be consulted for advice. Only the rectal route of administration should be adopted.
Do not exceed 3 days of treatment in event of fever or 5 days of treatment in the event of pain without the advice of doctor.

CAUTION:
Overdose may be injurious to liver.

STORAGE:
Store protected from light and moisture at a temperature not exceeding 25°C.
Keep the medicine out of reach of children

Manufactured by:



**AMERICAN
REMEDIES**
Care & Cure with Research
American Remedies Healthcare Pvt. Ltd.
(An ISO 9001-2015 Certified Company)

Gala No. 105 to 111, 1st Flr., Bldg. No. E-15/A,
Harihar Complex, Bhiwandi, Mumbai - 421302
At :334/5 GIDC Estate, Waghodia,
Dist. Vadodara, Gujarat, India-391760
E-mail: info@americanremedies.in
ICO: American Remedies LLC, DE, USA
© Copyright

Mfg. Lic. No.: G/25A/6163-A

135.00 mm