

being given too much Bupivacaine are usually as follows:

- Feeling dizzy or light-headed.
- Numbness of the lips and around the mouth.
- Numbness of the tongue.
- Hearing problems.

- Problems with your sight (vision).

To reduce the risk of serious side effects, your doctor will stop giving you Bupivacaine as soon as these signs appear. This means that if any of these happen to you, or you think you have received too much Bupivacaine, **tell your doctor immediately.**

More serious side effects from being given too much Bupivacaine include twitching of your muscles, fits (seizures), being sick (vomiting) and loss of consciousness.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects have also been reported.

Severe allergic reactions (rare, may affect up to 1 in 1,000 people)

If you have a severe allergic reaction, tell your doctor immediately. The signs may include sudden onset of:

- Swelling of your face, lips, tongue or throat. This may make it difficult to swallow.
- Severe or sudden swelling of your hands, feet and ankles.
- Difficulty breathing.

- Severe itching of the skin (with raised lumps).

Other possible side effects:

Very common (may affect more than 1 in 10 people)

- Low blood pressure. This might make you feel dizzy or light-headed.

- Feeling sick (nausea).

Common (may affect up to 1 in 10 people)

- Being sick (vomiting).

- Feeling dizzy.

The following information is intended for healthcare professionals only:

Any unused solution should be adequately disposed of, in accordance with local requirements.

1. Administration

Solution for injection.

The medicinal product is for epidural use, intraarticular use, subcutaneous use or perineural use only. The

maximum dosage must be determined by evaluating the size and physical status of the patient and considering the usual rate of systemic absorption from a particular injection site.

Experience to date indicates a single dose of up to 150 mg

bupivacaine hydrochloride monohydrate. Doses of up to 50 mg 2-hourly may subsequently be used. A maximum

dose of 2 mg/kg should not be exceeded in any four-hour

period. For young, elderly or debilitated patients, these doses should be reduced.

2. Handling Instructions

For single use only.

Only clear solutions practically free from particles should be used. Any unused solution should be discarded.

Do not use Bupivacaine Injection after the expiry date as indicated on the box after "EXP". The expiry date refers

to the last day of that month.

- Pins and needles.

- High blood pressure (hypertension).

- Slow heart beat.

- Problems passing water.

Uncommon (may affect up to 1 in 100 people)

- Feeling light-headed.

- Fits (seizures).

- Numbness of the tongue or around the mouth.

- Ringing in the ears or being sensitive to sound.

- Difficulty in speaking.

- Blurred sight or double vision.

- Loss of consciousness.

- Shaking (tremors).

- Twitching of your muscles.

Rare (may affect up to 1 in 1,000 people)

- Nerve damage that may cause changes in sensation or muscle weakness (neuropathy). This may include peripheral nerve damage.

- A condition called arachnoiditis (inflammation of the membrane that surrounds the spinal cord). The signs

include a stinging or burning pain in the lower back or legs and tingling,

numbness or weakness in the legs.

- Weak or paralysed legs.

- Uneven heart beat (arrhythmias). This could be life-threatening.

- Slowed or stopped breathing or stopped heart beat. This could be life-threatening.

Possible side effects seen with other local anaesthetics which might also be caused by Bupivacaine

include:

- Damaged nerves. Rarely this may cause permanent problems.

- Blindness which is not permanent or problems with the muscles of the eyes that are long-lasting. This may

happen with some injections given around the eyes.

Additional side effects in children and adolescents

Side effects in children are similar to those in adults.

Do not be concerned by this list of possible side effects. You may not get any of them.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in

this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine

5. How to store Bupivacaine

medicine : Keep out of reach of children.

Do not use this medicine after the expiry date which is stated on the vial and the box after "EXP". The expiry date

refers to the last day of that month.

Store below 25°C. Protected from light & moisture.

Shelf life after dilution:

From a microbiological point of view the product should be used immediately.

After first opening: to be used immediately

Only clear solutions practically free from particles should be used. Do not use if container is damaged.

Any unused solution should be discarded.

Do not throw away any medicines via waste-water or with household waste. Ask your pharmacist how to throw

away medicines you no longer use. These measures will help to

protect the environment.

What Bupivacaine looks like and content of the pack:

Bupivacaine is a clear, colourless, aqueous, sterile solution for injection supplied in 20 ml glass vial.

Manufactured by:



Call No. 105 to 111, 4th Fl. Bldg. No. E-15A,
Harhar Complex, Bhandari, Mumbai - 421302
At: VII, Surajpur Nahar Road Pancha Sahib,
Dist. Simlaur, Himachal Pradesh-173025.
E-mail: info@americanremedies.in

© Copyright



For the use only of a Registered Medical Practitioner of a Hospital or a Laboratory

Bupivacaine Hydrochloride Injection IP 100mg/20ml

AMERICAN
SENSOROX 0.5%
Injection

Local Anaesthetic

Composition :

Each ml contains :

Bupivacaine Hydrochloride IP 5mg

Sodium Chloride IP 8mg

Water for Injection IP q.s.

Read all of this leaflet carefully before you start using this medicine because it contains important information for

you.

- If you have any further questions, ask your doctor or pharmacist or nurse.

- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not

listed in this leaflet. See section 4.

What is in this leaflet

2. What you need to know before you use Bupivacaine

- Keep this leaflet. You may need to read it again.

3. How to use Bupivacaine

4. Possible side effects

1. What Bupivacaine is and what it is used for

5. How to store Bupivacaine

6. Contents of the pack and other information

1. What Bupivacaine is and what it is used for

Bupivacaine contains the active substance Bupivacaine hydrochloride monohydrate

it belongs to a group of medicines called local anaesthetics.

Bupivacaine is used to numb (anaesthetise) parts of the body. It is used to stop pain happening or to provide pain

relief. It can be used to:

- Numb parts of the body during surgery in adults and children above 12 years.

- Relieve pain in adults, infants and children above 1 year of age

2. What you need to know before you use Bupivacaine

Do not use Bupivacaine:

- If you are allergic to bupivacaine hydrochloride monohydrate, or any of the other ingredients of this medicine

- If you are allergic to any other local anaesthetics of the same class (such as lidocaine or ropivacaine).

- If you have a skin infection near to where the injection will be given.

You must not be given this medicine if any of the above apply to you. If you are not sure, talk to your doctor or

nurse before you are given this medicine.

Warnings and precautions

Talk to your doctor or pharmacist or nurse before using Bupivacaine

- If you have heart, liver or kidney problems. This is because your doctor may need to adjust the dose of

Bupivacaine. Problems with your liver enzymes may occur, particularly if

you have long-term treatment with this medicine.

- If you have epilepsy.

- If you have been told that you have decreased volume of blood (hypovolaemia).

Children

In children aged less than 12 years as some injections of Bupivacaine in order to numb parts of the body during

surgery are not established in younger children. The use of Bupivacaine is not established in children less than 1

year of age.

If you are not sure if any of the above apply to you, talk to your doctor or nurse before you are given this

medicine.

Other medicines and Bupivacaine

Tell your doctor or pharmacist if you are taking have recently taken or might take any other medicines. This

includes medicines that you buy without a prescription and herbal medicines. This is because Bupivacaine can

affect the way some medicines work and some medicines can have an effect on Bupivacaine.

In particular, tell your doctor if you are taking any of the following medicines:

- Medicines used to treat an uneven heart beat (arrhythmia) such as lidocaine, mexiletine or amiodarone.

Your doctor needs to know about these medicines to be able to work out the correct dose of Bupivacaine for you

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor

or pharmacist for advice before this medicine

Pregnancy

There are no or limited amount of data from the use of bupivacaine in pregnant women.

Breast-feeding

Bupivacaine enters the mother's milk, if you are breast-feeding you should discuss options with your doctor

Fertility

There are no data on the effect of bupivacaine on human fertility

Driving and using machines

Bupivacaine may make you feel sleepy and affect the speed of your reactions. After you have been given

Bupivacaine, you should not drive or use tools or machines until the next

day.

Bupivacaine Solution contains sodium

4. Possible side effects

The following side effects have also been reported:

This medicinal product contains 3.15 mg/ml (0.14 mmol) sodium per dose. Your doctor will take this into account if

you are on a sodium controlled diet.

3. How to use Bupivacaine

Bupivacaine will be given to you by your doctor. Your doctor will know the correct way to give you this medicine.

The dose that your doctor gives you will depend on the type of pain relief that you need and the part of your body

that the medicine will be injected into. It will also depend on your body size, age, and physical condition. Usually

one dose will last long enough but more doses may be given if the surgery takes a long time.

Bupivacaine will be given to you as an injection or infusion. The part of the body where you are injected will

depend on why you are being given Bupivacaine. Your doctor will give you

Bupivacaine in one of the following places:

- Near to the part of the body that needs to be numbed

In an area away from the part of the body that needs to be numbed. This is the case if you are given an epidural

injection (an injection around the spinal cord).

When Bupivacaine is injected into the body in one of these ways, it stops the nerves from being able to pass pain

messages to the brain. It will slowly wear off when the medical procedure is over.

Use in children and adolescents

Depending on the type of required analgesia Bupivacaine is injected slowly either into the epidural space (part of

the spine) or other parts of the body by a doctor experienced in

paediatric anaesthetic techniques. Dosage depends on the age and weight of the patient and will be determined

by the doctor.

If you use more Bupivacaine than you should

Serious side effects from getting too much Bupivacaine are unlikely. They need special treatment and the doctor

treating you is trained to deal with these situations. The first signs of